

Program and abstracts. JSLOC2024: 2nd JAPANESE-SPANISH SYMPOSIUM ON LOCALIZATION AND NONLINEAR PHENOMENA IN LATTICES AND WAVES, Grau de Gandia, Spain, March 4-9, 2024



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

PROGRAM AND ABSTRACTS

Premises: Escola Politècnica Superior de Gandia, Carrer del Paranimf, 1, 46730 Gandia, València

URL: <https://grupo.us.es/gfnl/JSLoc2024/>

Organizers: JFR Archilla (Universidad de Sevilla), Masayuki Kimura (Setsunan University), Víctor J- Sánchez-Morcillo (Universitat Politècnica de València) and Yusuke Doi (Osaka University).

PROGRAM:

Monday, 04/03/2024

Welcome meeting: 17:30-19:30:

Laboratory of Applied Physics: Experiments with breathers and other localized excitations. Research at the Laboratory of the Group of Applied Physics (UPV). Chair: VJ Sánchez-Morcillo.

Visit to Grau de Gandia and dinner at Barracuda restaurant.

Tuesday 05/03/2024

Opening: 10:00: Juan FR Archilla

Project Presentations Session. Chair: : v] • i Œ •

10:30 Muriel Botey/Ramón Herrero

Exploring complex spacetime systems for unconventional light control

11:00: Víctor Sánchez-Morcillo/Ruben Picó

Exploring complex spacetime systems for unconventional sound control

11:30 Coffe break

Lecture Session 1. Chair: Ruben Picó

12:00 Yusuke Doi, Osaka University

Dynamics of waves in pairwise interaction symmetric lattice

12:30 Yosuke Watanabe, Setsunan University

Measuring velocity of supratransmission in mass-spring chain

13:00 Masayuki Kimura, Setsunan University

Experiments on localized vibrations in magneto-mechanical resonators arrays

Lunch at the university restaurant

15:00 Laboratory visit and discussions: Exploring complex spacetime systems for unconventional sound control. **Chair: Ruben Picó**

17:00 Visit to Ducal Palace in Gandia

18:00 Guided visit to Gandia city

20:30 Conference Dinner at Restaurant Vines i Mes in Gandia



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Wednesday 06/03/2024

Lecture Session 2. Chair: Masayuki Kimura.

9:00 Juan FR Archilla, Universidad de Sevilla

Nonlinear excitations in a silicate chain. Experiments and theory

9:30 J. V. J. , University of Latvia

Efficient thermostating of semiclassical Hamiltonian lattice dynamics

10:00 Kestutis Staliunas, Universitat Politècnica de Catalunya

Hermitian and Non-Hermitian Light Propagation Effects in Thin Films

10:30 Ramón Herrero, Universitat Politècnica de Catalunya

Stabilization of laser and laser array emission by symmetric and asymmetric couplings from non-Hermitian potentials

11:00 Coffe break

Lecture Session 3. Chair: Muriel Botey.

11:30 Rubén Picó, Universitat Politècnica de València

Time-Varying and Dynamic Acoustic Metamaterials

12:00 Víctor Sánchez-Morcillo, Universitat Politècnica de València

Localized waves in highly nonlinear lattices: theory and experiments.

PhD Student Lecture Session. Chair: Muriel Botey.

12:30 Salim B Ivars, Universitat Politècnica de Catalunya

Hybridisation of the different scenarios leading to spatial localisation: hybrid frequency combs in non-Hermitian Kerr Cavities.

12:50 Nayeem Akhter, Universitat Politècnica de Catalunya

Unidirectional mode coupling in non-Hermitian waveguides

13:10 Jaime Galiana, Universitat Politècnica de València

Simulation methods for the study of time-varying acoustic systems.

13:30 Lunch: Fideua at Hogar del Pescador within the port of Grau de Gandia

15:30 Discussions: Exploring complex spacetime systems for unconventional sound control

17:30 Lighthouse walk. Dinner at No Name in Grau de Gandia.

Thursday 07/03/2024

10:00 Discussion panel 1: Breathers in molecular dynamics. Chair: Yusuke Doi

12:00 Discussion panel 2: Experiments on breathers: Chair: Yosuke Watanabe

Lunch:

15:00: Private discussions

Cultural activities: Guided visit to Real Monasterio de San Jeroni de Cotalba.

Friday 08/03/2024. Molecular Science Institute (ICMOL). Burjassot Campus. University of València.

11:00 Modulation of the optical properties of $\text{CrI}_3=\text{CrI}_3(\text{n-L})$. By Andrés Cantarero

12:00 The Laboratory of optical properties of two dimensional semiconductors. By Andrés Cantarero.

14:30 Lunch: Paella at Levante Restaurant in Benisanó.

16:30 Visit to València guided by Profs. Andrés Cantarero and Lluís M García-Raffi.

Saturday 09/03/2024

Discussion Panel 3: Breathers in electrical lattices. Chair: M Kimura

Closing: Juan FR Archilla



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

BOOK OF ABSTRACTS

JSLoc2024: 2nd Japanese-Spanish Symposium on Localization and Nonlinear Phenomena in Lattices and Waves

Grau de Gandia, Val+encia, Spain, March 4-9, 2024

Premises: Escola Politècnica Superior de Gandia, Carrer del Paranimf, 1, 46730 Gandia, València

URL: <https://grupo.us.es/gfni/JSLoc2024/>

Organizers:

Juan FR Archilla

Universidad de Sevilla, Sevilla, Spain

Masayuki Kimura

Setsunan University, Osaka, Japan

Víctor J Sánchez-Morcillo

Universitat Politècnica de València, Gandía, Spain

Yusuke Doi

Osaka University, Osaka, Japan



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Nonlinear excitations in a silicate chain. Experiments and theory.

Juan FR Archilla¹, Yusei Doi³, Masayuki Kimura⁴

¹Universidad de Sevilla, Sevilla, Spain; ²University of Latvia, Riga, Latvia;

³Osaka University, Osaka, Japan; ⁴Setsunan University, Osaka, Japan.

We present some experiments in silicates that show the existence of nonlinear excitations that transport energy and can transport electric charge without the presence of an electric field. The energy of the nonlinear excitations is provided by the impact of alpha particles. We present different models for which it is possible to obtain long-



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Efficient thermostating of semiclassical Hamiltonian lattice dynamics

J. Archilla¹, J. Bajars², Y. Doi², M. Kimura², F.R. Archilla²

¹University of Latvia, Riga, Latvia; ²Universidad de Sevilla, Sevilla, Spain

In this work, we propose mixed canonical-microcanonical equilibrium distribution and develop thermostat methods for semiclassical Hamiltonian lattice equations. In semiclassical Hamiltonian lattice models, the crystal lattice is described by classical Hamiltonian dynamics, whereas an extra charge (electron or hole) is modeled as a quantum particle within the tight-binding approximation. Such models are of significant scientific importance. A particular application is hyperconductivity, i.e., the experimental observation of charge transport without the presence of an external electric field when a silicate is bombarded with alpha particles. The charge is carried through the crystal by nonlinear lattice excitations. In the present work, the canonical equations for a semiclassical Hamiltonian describing the coupled lattice-charge dynamics are coupled to an efficient stochastic thermostat, which drives the system to the equilibrium distribution at a prescribed temperature with minimal perturbations to the Hamiltonian trajectories while at the same time ensuring the conservation of the charge probability. The properties of the proposed efficient thermostating are explored and numerically demonstrated on a phenomenological semiclassical Hamiltonian lattice model.

References

- [1] JFR Archilla, J Bajars, Y Doi and M Kimura. A semiclassical model for charge transfer along ion chains in silicates. J. Phys: Conf. Ser. (to appear), arXiv:2308.1518 (2024).
- [2] JFR Archilla and J Bajars. Spectral properties of exact polarobreathers in semiclassical systems. Axioms 12, 5 (2023) 437/1-26.
- [3] J Bajars and JFR Archilla. Splitting methods for semi-classical Hamiltonian dynamics of charge transfer in nonlinear lattices. Mathematics 10, 19 (2022) 3460.
- [4] FM Russell, MW Russell and JFR Archilla. Hyperconductivity in fluorphlogopite at 300 K and 1.1 T. EPL 127, 1 (2019) 16001.

Acknowledgments

The authors acknowledge the following projects and grants:

Project from the Faculty of Physics, Mathematics and Optometry, University of Latvia (2024); MICINN PID2019-109175GB-C22, MICINN PID2022-138321NB-C22, and travel grants from VII PPITUS-2024 of the Universidad de Sevilla.



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Exploring complex spacetime systems for unconventional light and sound wave control (STWaveControl)

Muriel Botey¹, Ramon Herrero¹, Salim B. Ivars¹, Nayeem Akhter¹, Kestutis Staliunas^{1,2}.

¹Universitat Politècnica de Catalunya, ²ICREA

We present and overview of the current state and ongoing works of the subproject from Universitat Politècnica de Catalunya (UPC) of the coordinated project Exploring complex spacetime systems for unconventional light and sound wave control (STWaveControl) [1,2,3].




Fig.1. a) Consecutive generations in metamaterial evolution [4] b) Generalized Hilbert Transform to design non-Hermitian media to control the field flow, corresponding to the 3rd metamaterials generation c) Multicolor-HT to design frequency selective materials

The present project aims at studying complex artificial metamaterials including spacetime structures for the control of waves, see Fig. 1. The scope of the project holds an intrinsic multidisciplinary character since effects arising from the interaction between waves and artificial spacetime structured materials apply generally to light and sound waves. Such effects form the basis of the present proposal for a collaboration between the Nonlinear Dynamics, Nonlinear Optics and Lasers research group at the UPC and the Acoustics Waves in Complex Media research group at the Universitat Politècnica de València.

References

- [1] SB Ivars, M. Botey, R. Herrero, K. Staliunas, K. (2023). Stabilisation of spatially periodic states by non-Hermitian potentials. *Chaos, Solitons & Fractals*, 168, 113089.
- [2] MN Akhter, M. Botey, R. Herrero, K. Staliunas (2024). Mode-cleaning in antisymmetrically modulated non-Hermitian waveguides. *Nanophotonics*.
- [3] SB Ivars, M. Botey, R. Herrero, K. Staliunas. Hybrid patterns and solitonic frequency combs in non-Hermitian Kerr Cavities, submitted
- [4] C Caloz, ZL Deck-Léger (2019). Spacetime metamaterials—part I: general concepts. *IEEE Transactions on Antennas and Propagation*, 68(3), 1569-1582.

Acknowledgments: The authors acknowledge the project MICINN PID2022-138321NB-C21.



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY


Dynamics of waves in a pairwise interaction symmetric lattice.

Yusuke Doi


Osaka University, Osaka, Japan

The pairwise interaction symmetric lattice (PISL) is a nonlinear lattice that supports smooth mobility of discrete breathers (DBs) by introducing long-range nonlinear interactions satisfying a certain symmetry of interactions[1, 2]. In this talk, we introduce the structure of the PISL. Moreover, some numerical results of dynamics of waves in the PISL such as modulational instability of zone boundary mode and interaction between moving DBs and normal modes are presented.


$T = 2.0, \nu = 1/10$




$T = 2.0, \nu = 1/5$



$T = 2.0, \nu = 1/3$



$T = 2.0, \nu = 1/1$



References:

- [1] Y Doi, K Yoshimura, Phys. Rev. Lett., Symmetric potential lattice and smooth propagation of tail-free discrete breathers, Phys. Rev. Lett., 117, 014101 (2016).
- [2] Y Doi, K Yoshimura, Construction of symmetric lattice for smooth mobility of discrete breathers, Nonlinearity, 33, 5142 (2020).



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505




大阪大学
OSAKA UNIVERSITY

Experiments on Spatially Localized Vibrations in Magneto-mechanical Resonator Array

Masayuki Kimura

Setsunan University, Osaka, Japan

Spatially localized vibrations in nonlinear resonator arrays, which are called intrinsic localized modes (ILMs) or discrete breathers (DBs), have been observed in a variety of real physical systems such as, even in mechanical systems, nonlinearly coupled pendula [1,2], micro-mechanical cantilever arrays [3], and linearly coupled pendula [4]. We have also fabricated magneto-mechanical resonator arrays [5,6] to observe ILM/DB and to manipulate its position. The photos of the arrays are shown below. Recently, we have studied magnetically coupled two-degree-of-freedom resonators. In this presentation, experimental systems of the magneto-mechanical resonator arrays will be introduced with observation results, and recent progress on the two-degree-of-freedom resonator arrays will be shown.



Photos of the magneto-mechanical resonator arrays. Left: Linearly coupled cantilever array [5]. Right: Magnetically coupled cantilever array [6].

References

- [1] FM Russell, Y Zolotaryuk, JC Eilbeck, and T Dauxois, Moving breathers in a chain of magnetic pendulums, *Phys. Rev. B* 55(10), 6304 (1997).
- [2] Y Watanabe, T Nishida, Y Doi, and N Sugimoto, Experimental demonstration of excitation and propagation of intrinsic localized modes in a mass-spring chain, *Phys. Lett. A* 382 (30), 1957 (2018).
- [3] M Sato, BE Hubbard, AJ Sievers, B Ilic, DA Czaplewski, and HG Craighead, Observation of Locked Intrinsic Localized Vibrational Modes in a Micromechanical Oscillator Array, *Phys. Rev. Lett.*, 90, 44102 (2003).
- [4] J Cuevas, LQ English, PG Kevrekidis, and M Anderson, Discrete Breathers in a Forced-Damped Array of Coupled Pendula: Modeling, Computation, and Experiment, *Phys. Rev. Lett.*, 102, 224101 (2009).
- [5] M Kimura and T Hikiara, Coupled Cantilever Array with Tunable On-site Nonlinearity and Observation of Localized Oscillations, *Phys. Lett. A* 373, 1257 (2009).
- [6] M Kimura, Y Matsushita, and T Hikiara, A Study on Bifurcations and Structure of Phase Space Concerning Intrinsic Localized Modes in a Nonlinear Magneto-Mechanical Lattice, *AIP Conference Proceedings* 1474 (ISNA19), 55-58 (2012).

Acknowledgments: The beginning of this work was supervised by Prof. T. Hikiara (Kyoto University, Japan). This work is supported by JSPS Kakenhi No. 22760280, 25820164, 18K04020, and 21K03935. The author also acknowledges travel support from project MICINN PID2022-138321NB-C22.



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Stabilization of laser and laser array emission by symmetric and asymmetric couplings from non Hermitian potentials

Ramon Herrero

Universitat Politècnica de Catalunya, Spain

Spatiotemporal instabilities in lasers and laser arrays induce temporally unstable and low spatial quality emission. Simultaneous intracavity modulations of refractive index and gain-loss lead to efficient stabilization mechanisms. The stabilization can be based on the suppression of the modulation instability, on the stabilization of specific stationary solutions or on field localization. The stabilization mechanism is shown in Vertical Cavity Semiconductor Emitting Lasers, Edge Emitting Lasers and arrays of Edge Emittig Laser.



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Time-Varying and Dynamic Acoustic Metamaterials

Rubén Picó¹, Javier Redondo¹, Víctor J. Sánchez-Morcillo¹.

¹Instituto de Investigación para la Gestión Integrada de Zonas Costeras, Universitat Politècnica de València, Campus de Gandía.

Recent advances in metamaterials have dramatically extended the range of modern metamaterial properties [1]. Time-varying metamaterials are materials whose properties depend on the temporal variation of their physical parameters induced by an external source of energy [2]. We propose a



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Localized waves in highly nonlinear lattices: theory and experiments

Víctor J Sánchez-Morcillo¹, Vitaliy Gusev², Noé Jiménez³, and Rubén Picó¹

³ Instituto de Investigación para la Gestión Integrada de Zonas Costeras (IGIC), Universitat Politècnica de València, Paraninf 1, 46730 Gandia (Spain), victorsm@upv.es

² Le Mans Université, av. O. Messiaen, 72085, Le Mans (France), vitali.goussev@univ-lemans.fr

³ Instituto de Instrumentación para Imagen Molecular (i3M), Universitat Politècnica de València, Camino de Vera S/N, 46020 València (Spain), nojigon@i3m.upv.es

Summary: Wave propagation in a highly nonlinear system of coupled rotators is investigated. A discrete nonlinear wave equation is obtained, and the main characteristics of its propagating solutions are discussed. An experiment was designed to test the analytical predictions. Nonlinear signatures as amplitude-dependent dispersion, bistability and self-modulations, and strong localization are observed and discussed.

Highly nonlinear systems, including those where any linear response is absent (fully nonlinear systems) have been much less investigated than weakly nonlinear systems. Some known examples are granular media [1], or connected discs performing torsional motions [2]. In such systems, any arbitrarily small perturbation may lead to significant nonlinear effects. The absence of linear waves defines also the so-called *acoustic vacuum*, a medium where the speed of sound as defined by classical acoustics vanishes. Such systems are also known to support highly localized waves or *compactons*.




Figure 1 Three neighbour elements of the infinite rotator lattice. a) The lattice at rest, with rotating elements connected by unstressed springs. b) Lattice excited with a symmetric mode $k=\pi$ and c) Lattice excited with the antisymmetric mode $k=2\pi$.

We present a theoretical/experimental study of the dynamics of mechanical waves propagating in a highly nonlinear one-dimensional lattice consisting of rotating elements, with mass m and inertia moment I , coupled by linear springs (Fig. 1). The nonlinearity, of geometrical origin, is of cubic type. In the case of absence of pre-stress of the springs in the rest state (Fig. 1a), linear terms are absent and the system is purely nonlinear. The equation of motion for this lattice has been derived for a general geometry (for arbitrary spring-to-rotator length ratio), and its different limits have been explored. The equation of motion for rotator n is defined by its angle φ_n . Wave propagation in this system is described by the nonlinear discrete equation

$$\mu \ddot{\varphi}_n + a \varphi_n^3 + b \varphi_{n-1} \varphi_n \varphi_{n+1} = 0 \quad (1)$$

where a , b and μ are geometrical parameters dependent on the spring length l_0 and radius of rotator R . As the system described by Eq. (1) is highly nonlinear, it does not show linear dispersion at all, presenting what has been called an acoustic vacuum. Moreover, the obtained nonlinear dispersion is of optical phonon type, i.e. the wavelength increases with the frequency of the wave. Figure 2 shows the dispersion relation of the first harmonic (the wave at the excitation frequency). Note the presence of upper-lower bandgaps. Note also that, since the frequency in Fig. 2 is normalized to the amplitude of



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

excitation, this means that, for a given excitation frequency, the dispersion relation can be tuned by the excitation amplitude.

Analytical solutions for the amplitude of the fundamental wave and its third harmonic have been obtained in the rotating wave approximation. Other interesting feature of the system is that the amplitude of the third harmonic vanishes for a particular k mode, i.e. the system supports purely harmonic waves.




Figure 2 Dispersion relation for the case $l_0=R$ ($\mu=0.5$). Frequency is normalized to the amplitude of the excitation, therefore waves can be propagating or evanescent depending on the initial amplitude.

Notably, we also obtain highly localized solutions (compacton-like), and study their existence conditions.

An experiment has been designed to check the main predictions of the theory. The system is excited at the boundary by driving harmonically the angle of the first rotator, and the motion of each rotator is tracked by video and processed to obtain the time evolution of the angles. Measurements show evidences of the nonlinear characteristics of the system, as the generation of harmonics, bistability and self-modulations, and the predicted amplitude-dependent transition from evanescent to propagating

References

- [1] V Nesterenko, Dynamics of heterogeneous materials, Springer (2001).
- [2] OV Rudenko and EV Solodov, *Acoust. Phys.* **57**, 51-58 (2011).
- [3] YS Kivshar, *Phys Rev. E* **48**, R43 (1993).
- [4] P Rosenau, *Phys. Rev. Lett.* **73**, 1737 (1994).

Acknowledgments: VSM acknowledge support from the projects PID2019-109175GB-C22 and PID2022-138321NB-C22



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505




大阪大学
OSAKA UNIVERSITY

Measuring velocity of supratransmission in a mass-spring chain


Yosuke Watanabe

Setsunan University, Osaka, Japan

Propagation velocity of nonlinear localized oscillations in a mass-spring chain is measured. The chain emulates the Fermi-Pasta-Ulam one of beta type and sinusoidal driving with large amplitude and high frequency at the one end may excite the propagating localized oscillations [1], the supratransmission [2,3]. The propagating velocity appears to be nearly constant throughout the propagation.



Snapshot of the propagation of nonlinear localized oscillation in the mass-spring chain. In the movie some oscillators are marked with color points to be tracked.



Tracking results of the propagations of localized oscillations, in which color coding coincides with the figure shown above.

References

- [1] Y Watanabe, et al., Phys. Lett. A 382 (2018) 1957-1961.
- [2] R Khomeriki, et al., Phys. Rev. E 70 (2004) 066626.
- [3] F Geniet and J Leon, Phys. Rev. Lett. 89, 13 (2002) 134102.

Acknowledgments

This work was supported by JSPS KAKENHI Grant Number 23K03582.



UNIVERSITAT
POLITÀCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

PHD STUDENTS SESSION:

Unidirectional mode coupling in non-Hermitian waveguides

Nayeem Akhter

Universitat Politècnica de Catalunya

We propose an all-optical control of the mode dynamics in multimode waveguides, by simultaneous modulation of the refractive index and of the gain/loss coefficient. The induced unidirectional mode-coupling, yields to the enhancement or reduction of the excitation of higher order transverse modes, depending on the modulation parameters. In the latter case, the all-optical mode-cleaning is observed, resulting in an almost single-mode spatially coherent output. While coupling towards higher order modes, enhances pulsing and, eventually, leads to supercontinuum generation. We analytically predict the effect on a coupled mode theory for 1D waveguides, which is numerically proven solving the wave propagation equation. The proposal is generalized to the more involved case of 2D waveguides for different geometries. We expect these results to pave the way towards novel non-Hermitian photonics applications.

Hybridisation of the different scenarios leading to spatial localisation: hybrid frequency combs in non-Hermitian Kerr Cavities.

Salim Benadouda Ivars

This study introduces a novel scenario for the formation of dissipative localised structures in nonlinear systems. Traditionally, soliton formation relies on the existence of at least two different potentially stable stationary states: either two uniform states (flat), or a uniform and a patterned state. Each formation scenario holds unique features, which can be typically found in optical cavities with either normal and anomalous dispersion, respectively. However, we show how the introduction of a periodic non-Hermitian modulation in Kerr cavities induces a hybridisation of these two established mechanisms, resulting in a new scenario that combines properties of both. As a consequence, the basal stationary states exhibit dual behavior, where a state can act as either flat or patterned under the same parameters. This phenomenon arises from the stabilization of periodic solutions comprising trains of dark solitons, which are unstable in the unmodulated case.

These findings expand the fundamental understanding of localised structures in nonlinear science, introducing a new paradigm for their formation. Beyond a fundamental scope, the work holds significant implications, especially for frequency comb generation, as the hybridisation blends anomalous and normal dispersion properties within a normal dispersive cavity. This allows for the stabilisation of new families of frequency combs associated to stable solitons, molecules, and patterns. Additionally, it leads to an unexpected and reversible mechanisms for real-time comb manipulation and control.



UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA

Smart and Human
摂南大学



UNIVERSIDAD
DE SEVILLA
1505



大阪大学
OSAKA UNIVERSITY

Simulation methods for the study of time-varying acoustic systems

Jaime Galiana-Nieves¹, Javier Redondo¹, Víctor J. Sánchez-Morcillo¹, Rubén Picó¹.

¹Instituto de Investigación para la Gestión Integrada de Zonas Costeras, Universitat Politècnica de València, Campus de Gandía.


In the last years, the study of time-varying materials by means of simulations is gaining interest. In this work we focus on the FDTD method for the simulation of time-dependent models. Due to the complexity of implementing moving systems using this method, different techniques have been studied to approximate the solution of these systems.

Firstly, boundary conditions have been implemented as FIR filters obtained from the acoustic admittance calculated using the transfer matrix method. In this way, the response of the system at each time instant can be calculated and applied at a single point.

Secondly, FDTD has been combined with the Immersed Boundary Method (IBM) for the simulation of time-varying systems.

Finally, an implementation has been implemented from analytical expressions of a static Helmholtz resonator. Once the system has been validated, we will proceed to the implementation of the same system, but time-variant.

Figure: Dispersion relation of an static Helmholtz resonator obtained from the analytical expression (purple) and the results from the FDTD simulations (white).



References

- [1] D Zhao, CA Barrow, AS Morgans, and J Carrote, Acoustic Damping of a Helmholtz Resonator with an Oscillating Volume, *AIAA Journal*, 47, 7, 2009.
- [2] E Riva, MIN Rosa, Y Guo, and M Ruzzene, Adiabatic sound transport in acoustic waveguides with time-varying Helmholtz resonators, *Frontiers in Acoustics*, 2023.
- [3] AG Hayrapetyan, KK Grigoryan, RG Petrosyan, and S Fritzsche, Propagation of sound waves through a spatially homogeneous but smoothly time-dependent medium, *Ann. Phys.*, 333, 47-65, 2013